



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$m=(c-b)^2y$ .  $\therefore y = \frac{m}{(c-b)^2}$ . The value of the work which could be done at the age of  $c-x$  years is,  $x^2y = \frac{mx^2}{(c-b)^2}$ .

$\therefore$  the value of total amount of work done between the ages of  $b$  and  $c$  years is,  $\frac{m}{(c-b)^2} \int_0^{c-b} x^2 dx = \frac{1}{3}m(c-b)$ .

Adding this to value given in (1), we have,  $\frac{1}{3}m(c-b) + m(b-a) = d$ , since  $d$  = value of horse's work from  $a$  to  $c$  years of age.

$\therefore m = \frac{3d}{c-3a+2b}$ . For any age,  $b-n$ , between the ages of  $a$  and  $b$  years, we have, the value of horse,  $V$  = value of work done after that age  $= mn + \frac{1}{3}m(c-b) = \frac{3d}{c-3a+2b} [n + \frac{1}{3}(c-b)] \dots (A)$ .

For any age,  $c-n'$ , between the ages of  $b$  and  $c$  years, we have, value of horse,

$$V' = \frac{m}{(c-b)^2} \int_0^{n'} x^2 dx = \frac{mn'^3}{3(c-b)^2} = \frac{dn'^3}{(c-b)^2(c-3a+2b)} \dots (B).$$

Ex. Let us suppose that a horse is worth \$100 at five years of age, and that he begins to weaken at 18 years of age, becoming worthless at 20. Then,  $a=5$ ,  $b=18$ ,  $c=20$ , and  $d=100$ . Substituting (A) becomes,

$V = \frac{100(3n+2)}{41}$ , (A), and (B), becomes,  $V' = \frac{25n'^3}{41} \dots (B)$ . By substituting for  $n$  and  $n'$  their successive integral values, we construct the following table:

years	value	years	value	years	value
5	\$100	10	\$63.41	15	\$26.82
6	92.68	11	56.09	16	19.51
7	85.36	12	48.78	17	12.19
8	78.04	13	41.46	18	4.87
9	70.73	14	34.14	19	.60
				20	.00

Also solved by F. P. MATZ, and G. B. M. ZERR.

29. Suggested by MANSFIELD MERRIMAN, C. E., Ph. D., Professor of Civil Engineering, Lehigh University, South Bethlehem, Pennsylvania.

Solve neatly the equations:  $\frac{y(1+x^2)}{x(1+y^2)} = a \dots (1)$ , and  $\frac{y^2(1+x^2)}{x^2(1+y^2)} = b \dots (2)$ .

I. Solution by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

Write (1) and (2), respectively,

$$x + \frac{1}{x} = a \left( y + \frac{1}{y} \right) \dots (3), \text{ and } x^2 + \frac{1}{x^2} = b \left( y^2 + \frac{1}{y^2} \right) \dots (4).$$

The second power, and the fourth power, of (3) give respectively,

$$x^2 + \frac{1}{x^2} = a^2 \left( y + \frac{1}{y} \right)^2 - 2 \dots (5), \text{ and } x^4 + \frac{1}{x^4} + 4 \left( x^2 + \frac{1}{x^2} \right) + 6$$

$$= a^4 \left( y + \frac{1}{y} \right)^4 \dots (6). \quad \text{From (6), by means of (5), we have } x^4 + \frac{1}{x^4} + 4$$

$$\left[ a^2 \left( y + \frac{1}{y} \right)^2 - 2 \right] + 6 = a^4 \left( y + \frac{1}{y} \right)^4 \dots (7).$$

$$\text{From (4) and (7), } \left( y + \frac{1}{y} \right)^4 - \left( \frac{4a^2}{a^4 - b} \right) \left( y + \frac{1}{y} \right)^2 = - \frac{2}{a^4 - b} \dots (8).$$

$$\therefore y + \frac{1}{y} = \frac{2a^2 \pm \sqrt{2(a^4 - b)}}{a^4 - b} = v \dots (9).$$

From (9),  $y = \frac{1}{2} [\sqrt{u} \pm \sqrt{u-4}] \dots (10)$ . From (3), by means of (9), we deduce  $x = \frac{1}{2} [a\sqrt{u} \pm \sqrt{a^2 u - 4}] \dots (11)$ .

II. Solution by G. B. M. ZERR, A. M., Principal of High School, Staunton, Virginia, and P. S. BERG, Apple Creek, Ohio.

We may write the expressions as follows:

$$\frac{1+x^2}{x} = a \left( \frac{1+y^2}{y} \right), \text{ and } \frac{1+x^6}{x^4} = b \left( \frac{1+y^6}{y^4} \right), \text{ or } x + \frac{1}{x} = a \left( y + \frac{1}{y} \right), \text{ and } x^4 + \frac{1}{x^4}$$

$$= b \left( y^4 + \frac{1}{y^4} \right). \quad \text{Let } x + \frac{1}{x} = z, \text{ and } y + \frac{1}{y} = v.$$

$$\therefore z = av \dots (1), \quad z^4 - 4z^2 + 2 = b(v^4 - 4v^2 + 2) \dots (2).$$

Substitute (1) in (2), we get,  $a^4 v^4 - 4a^2 v^2 + 2 = bv^4 - 4bv^2 + 2b$ .

$$\therefore (a^4 - b)v^4 - 4(a^2 - b)v^2 = 2(b - 1).$$

$$\therefore v^2 = \frac{2(a^2 - b) \pm \sqrt{[2(a^2 - b)^2 + 2b(a^4 - 1)]}}{a^4 - b} = u \text{ suppose.}$$

$$\therefore v = \pm \sqrt{u}, \quad z = \pm a\sqrt{u}.$$

$$\therefore x + \frac{1}{x} = \pm a\sqrt{u}, \text{ or } x^2 \pm a\sqrt{ux} = -1, \text{ and } y + \frac{1}{y} = \pm \sqrt{u}, \text{ or } y^2 \pm \sqrt{uy} = -1$$

$\therefore x = \pm \frac{1}{2} (a\sqrt{u} \pm \sqrt{a^2 u - 4})$ ,  $y = \pm \frac{1}{2} (\sqrt{u} \pm \sqrt{u-4})$ , which are the correct results, and the same as given by the proposed in a previous number of the MONTHLY.

NOTE—A neat solution to No. 26, Algebra, was received from Jno. B. Fraught, A. B., Instructor in Mathematics, Indiana University, after copy had been sent to the Publishers.

## PROBLEMS.

29. Proposed by ARTHURAS MARTIN, LL. D., U. S. Coast and Geodetic Survey Office, Washington, D. C.